

Causes moderate eye irritation. Avoid contact with eyes or clothing. Wash thoroughly with soap and water after handling.

FIRST AID: If in eyes, remove contact lenses. Hold eye open and slowly rinse with water. If irritation persists for more than 5 minutes, continue rinsing and consult a physician or poison control center.

Part No. ST50-600

EPA Est. No. 71979-SC-001

EPA Reg. No. 8383-7

With disinfectant

Net contents: 0.12 oz. 5" x 8"

(12.7 x 20.3cm) towelette saturated

with disinfectant

• **BACTERICIDAL**

• **FUNGICIDAL**

• **VIRUCIDAL****

• **TUBERCULOCIDAL**

• **PREVENTS MOLD**

• **AND MILDEW**

• **ELIMINATES ODORS***

Phenol.....	1.56%
Sodium Phenate.....	0.06%
Inert Ingredients.....	98.38%
Total.....	100.00%

Keep Out of Reach of Children

HAZARD TO HUMANS AND DOMESTIC ANIMALS

PRECAUTIONARY STATEMENTS:

HAZARD TO HUMANS AND DOMESTIC ANIMALS

Keep Out of Reach of Children

Active Ingredients

Phenol..... 1.56%

Sodium Phenate..... 0.06%

Inert Ingredients..... 98.38%

Total..... 100.00%

Disinfectant Towelette



Continuous Residual Activity
up to 6 Months*

ALCOHOL
FREE

AN EFFECTIVE CLEANER, DISINFECTANT AND ACTIVE DEODORIZER FOR INSTRUMENTS, AND EQUIPMENT ON PRE-CLEANED, HARD, NON-POROUS SURFACES in health/hospital treatment and patient rooms, operating rooms, schools, embassies, hotels, restaurants, boats, planes, buses, and the home. Recommended for medical and dental equipment, beds, surgical carts, counter tops, mannequins, telephones, keyboards, furniture, wheelchairs, walkers, sinks, doors, light switches, linen hampers, shower stalls, toilet seats, trash receptacles, bedrails, and food service areas. Resists the hazards of cross-contamination.

PASSES AOAC EFFICACY STANDARDS FOR DISINFECTION IN THE PRESENCE OF 5% ORGANIC SOIL, FOR INSTITUTIONAL, HOSPITAL AND FARM PREMISE USE.

Virucidal:** Avian Influenza A (H9N2), Influenza A (Japan 305/57 Asian strain),

Cytomegalovirus, Coronavirus, Polio Type 1 Virus, Vaccinia, Canine parvovirus, Herpes simplex

types 1F and 2/G (oral, ocular and genital), HIV-1 (AIDS Virus). See Sporicidin Bulletin NO. 303.

Bactericidal: Staphylococcus aureus, Salmonella enterica, Pseudomonas aeruginosa,

Methicillin-resistant Staphylococcus aureus (MRSA), vancomycin resistant Enterococcus faecalis (VRE), E. coli, Streptococcus pyogenes, Staphylococcus salivarius, Mycobacterium

tuberculosis.

Fungicidal: Chaetomium globosum, Aspergillus niger, Penicillium variable, Trichophyton

mentagrophytes (Athlete's Foot fungus).

ACTIVE DEODORIZER*: Provides continuous residual bacteriostatic activity against

odor-causing bacteria for up to 6 months, if not removed from surfaces including

plastics, glass, vinyl, treated wood, metal, glazed porcelain, glazed tile and paint.

SPORICIDIN DISINFECTANT TOWELETTES are germicidal when used as directed for

disinfection. For more information, contact Sporicidin products.

DIRECTIONS FOR USE: Use as directed on label. A violation of U.S. Federal Law to use this product in a manner

inconsistent with its labeling.

DO NOT DILUTE OR MIX WITH OTHER CHEMICALS

PRECLEAN surfaces using Sporicidin Disinfectant Solution to remove soil and filth. Wipe dry

with paper towel, cloth or sponge.

TO DISINFECT AND DEODORIZE items which cannot be immersed, such as electrical panels,

cash registers, door knobs, handles, seats, dashboards and other "high touch" surfaces in and

around farm equipment, animal and poultry housing, barns, kennels, breeding pens,

trucks and vehicles. Thoroughly wet pre-cleaned surface with Sporicidin

towelette. Allow to remain wet for 10 minutes. Repeat application if necessary. Do not use on any

surfaces that are porous or absorbent.

TO KILL, CLEAN AND PREVENT MOLD AND MILDEW: Use procedure stated above. Repeat

application if new growth appears.

STORAGE AND DISPOSAL: Store at room temperature. Discard empty container in regular

trash or other recycling if available.

FOR HOSPITAL DISINFECTION: This product is not to be used as a terminal disinfectant/high level

disinfectant on any surface or instrument that it is introduced directly into the human body,

or into any body fluid. It is not to be used on any surface that is in direct contact with

contacts intact mucous membranes but which does not ordinarily generate the blood barrier

or otherwise enter normally sterile areas of the body. This product may be used to pre-clean or

decontaminate critical or semi-critical medical devices prior to sterilization or high level

disinfection.

Manufactured for Corixa, Inc., Sporicidin Division, Spartanburg, S.C. 29303
1-800-424-3733 • www.sporicidin.com